

laboratories. The survey will be funded in full by the Office of Surveillance, Epidemiology, and Laboratory Services (OSELs) of the Centers for Disease Control and Prevention (CDC). Influenza epidemics usually cause an average more than 200,000 hospitalizations and 36,000 deaths per year in the U.S. Respiratory illnesses caused by influenza viruses are not easily differentiated from other respiratory infections based solely on symptoms. Also influenza viruses may adversely affect different subpopulations. The effective use of rapid influenza diagnostic testing practices is an important component of the differential diagnosis of influenza-like-illness in both inpatient and outpatient treatment facilities. Test results are used for making decisions about antiviral vs. antibiotic use, and in making admission or discharge decisions. In many cases, rapid influenza tests are the only tests that can provide results while the patient is still present in the facility. Thus, the appropriate use of the tests, and interpretation of test results is critical to the treatment and control of influenza. More than a dozen rapid tests

have been approved by the U.S. Food and Drug Administration and are in widespread use. The reliability of rapid influenza tests is influenced by the individual test product used and the setting. Reported sensitivities range from 10–75%; while the median specificities reported are 90–95%. Other factors influencing accuracy are the stage (or duration) of illness when the diagnostic specimen is collected, type and adequacy of the specimen collected, variability in user technique for specimen collection or assay performance, and disease activity in the community. Given these and other collective findings, it is imperative for public health and for response planning that CDC develops sector-specific guidance and effective outreach to the clinicians on appropriate use of RIDT in their practices.

Previous studies by CDC of outpatient facilities showed that clinical laboratories usually perform the rapid tests for emergency departments, and provide results for both inpatient and outpatient treatment. Thus, understanding the use of rapid influenza testing in clinical laboratories, how the laboratories report results to

emergency departments and treatment facilities and health departments, and what quality assurance practices are used will guide future efforts of the CDC to develop appropriate influenza testing guidelines and sector-specific training materials for clinicians and improve health outcomes of the American public.

The survey covers basic laboratory demographic characteristics, specimen collection and processing, testing practices, reporting of results to emergency departments and other treatment facilities, reporting results to health departments, quality assurance practices, and methods of receiving updated influenza-related information. The majority of the questions request information about laboratory influenza testing practices.

To date, no systematic study has been conducted to investigate how laboratories use these tests, how they report results, or how they interact with outpatient treatment facilities. The survey will be conducted on a national sample of clinical laboratories. There are no costs to respondents except their time.

ESTIMATED ANNUALIZED BURDEN HOURS

Type of respondents	Form name	No. of respondents	No. of responses per respondent	Avg. burden per response (in hrs)	Total burden (in hrs)
Clinical Laboratory Supervisors .....	Survey of Rapid Influenza Diagnostic Test Practices in Clinical Laboratories.	600	1	30/60	300
Total .....	.....	.....	.....	.....	300

Dated: June 1, 2011.

**Daniel Holcomb,**

*Reports Clearance Officer, Centers for Disease Control and Prevention.*

[FR Doc. 2011-14147 Filed 6-7-11; 8:45 am]

**BILLING CODE 4163-18-P**

**DEPARTMENT OF HEALTH AND HUMAN SERVICES**

**Centers for Disease Control and Prevention**

**Disease, Disability, and Injury Prevention and Control Special Emphasis Panel (SEP): Initial Review**

The meeting announced below concerns Strategies to Improve Vaccination Coverage of Children in Child Care Centers (CCCs) and Preschools, Funding Opportunity Announcement (FOA) IP11-006; Strategies to Increase Health Care

Providers Use of Population-Based Immunization Information Systems, FOA IP11-008; Effectiveness in an Intervention to Promote a Targeted Vaccination program in the Obstetrician-Gynecologist Setting, FOA IP11-009; initial review.

*Correction:* The notice was published in the **Federal Register** on April 29, 2011, Volume 76, Number 83, Pages 24031. The place should read as follows:

*Place:* Holiday Inn Decatur Conference Center, 130 Clairemont Avenue, Decatur, Georgia 30030, *Telephone:* (404)371-0204.

*Contact Person for More Information:* Gregory Anderson, M.S., M.P.H., Scientific Review Officer, CDC, 1600 Clifton Road, NE., Mailstop E00, Atlanta, Georgia 30333, *Telephone:* (404) 498-2293.

The Director, Management Analysis and Services Office, has been delegated the authority to sign **Federal Register**

notices pertaining to announcements of meetings and other committee management activities, for both the Centers for Disease Control and Prevention and the Agency for Toxic Substances and Disease Registry.

Dated: May 27, 2011.

**Elaine L. Baker,**

*Director, Management Analysis and Services Office, Centers for Disease Control and Prevention.*

[FR Doc. 2011-14202 Filed 6-7-11; 8:45 am]

**BILLING CODE 4163-18-P**